

Méthodes de lutte alternatives aux pesticides conventionnels contre les principaux ennemis du fraisier -

Catherine Thireau et Marianne Lefebvre, PRISME

Projet Prime-Vert : PRIS-2-16-1796

Durée : Février 2017/Mars 2018

Références bibliographiques par ravageur

- Anthonome du fraisier (*Anthonomus signatus*)

Baroffio, C., Borg-Karlsson, A., Cross, J., Fountain, M., Guibert, V., Hall, D., Ralle, B., Richoz, P., Rogivue, A., Sigsgaard, L., Trandem, N., Wibe, A., 2015. Comment attire deux ravageurs des framboises dans un même piège ? Revue suisse Viticulture, Arboriculture, Horticulture, vol. 47(3) :152-158.

Sabbahi, R., Merzouki, A., Guertin, C., 2008. Efficacy of Beauvaria bassiana against the strawberry parts, *Lygus lineolaris*, *Anthonomus signatus* and *Otiorhynchus ovatus*, *J. Appl. Entomol.* 132 151-160

Wibe, A., Borg-Karlsson, A.-K., Cross, J., Bichão, H., Fountain, M., Libikas, I., Sigsgaard, L., 2014. Combining 1,4-dimethoxybenzene, the major flower volatile of wild strawberry *Fragaria vesca*, with the aggregation pheromone of the strawberry blossom weevil *Anthonomus rubi* improves attraction. *Crop Protection* 64 (2014) 122-128.

- Punaise terne (*Lygus lineolaris*)

Agriculture et Agroalimentaire Canada, 2011. Stratégie de remplacement de lutte contre la punaise terne dans les fraiseraies de l'Ontario. No AAC 11419F; No de catalogue A118-43/2011F-PDF

Crampton, LA, Loeb, G., Hoelmer, K., Hoffmann, M., 2010. Effect of insecticide regimens on biological control of the tarnished plant bug, *Lygus lineolaris*, by *Peristenus* spp. in New-York state apple orchards. *Journal of Insect science*: volume 10

Crampton, LA. Biological control of the tarnished plant bug, *Lygus lineolaris* (Hemiptera: Miridae), by *Peristenus* spp. (Hymenoptera: Braconidae), in New York apple orchards. Masters Thesis, Cornell University; 2007.

Day WH, Hedlund RC, Saunders LB, Coutinot D, 1990. Establishment of *Peristenus digoneuti* (Hymenoptera: Braconidae), a parasite of the tarnished plant bug (Hemiptera: Miridae), in the United States. *Environ. Entomol.* 19, 1528. 1533.

Dumont, F. et Provost, C. 2016. Combinaison de méthodes de lutte pour contrer la punaise terne en fraiserie. Congrès conjoint de la Société de pédontomologie du Québec et de la Société de protection des plantes du Québec, Nicolet, 2-4 novembre 2016.

Easterbrook, M.A., Tooley, J.A., 1999. Assessment of trap plants to regulate numbers of the European tarnished plant bug, *Lygus rugulipennis*, on late-season strawberries. *Entomologia experimentalis et applicata*, volume 92, issue 2, 119-125.

Guertin, C., Sabbahi, R., Trudel, R., Jobin, É., 2004. Utilisation du champignon entomopathogène *Beauveria bassiana* contre les ravageurs des fraises. INRS. http://www.mapaq.gouv.qc.ca/SiteCollectionDocuments/Recherche_Innovation/Petitsfruits/Fiche102075.pdf

Loeb, G., Pritts, M., 2010. A trap crop system for managing tarnished plant bug damage in strawberries, Cornell University, New York State IPM Program

Sabbahi, R., Merzouki, A., Guertin, C., 2008. Efficacy of Beauvaria bassiana against the strawberry parts, *Lygus linolaris*, *Anthonomus signatus* and *Otiorhynchus ovatus*, *J. Appl. Entomol.* 132 151-160

Shelton, A. Biological Control: A guide to natural enemies in North America, Cornell University. <https://biocontrol.entomology.cornell.edu/index.php>

Swezey, S.L., Nieto, D.J., Pickett, C.H., Hagler, J.R., Bryer, J.A., Machtley, S.A. 2014. Spatial density and movement of the *Lygus* spp. Parasitoid *Peristonus relictus* (Hymenoptera: Braconidae) in organic strawberries with Alfalfa trap crops, *Entomological Society of America, Environ. Entomol.* 43(2): 363-369

Villeneuve, S., Bouchard, A., Rodrigue, F., 2011. Contrôle mécanique de la punaise terne dans la culture de fraise sur rangs nattés en régie biologique à l'aide d'une faucheuse rotative modifiée. Nord-Bio

<https://www.agrireseau.net/references/9/Journ%C3%A9e%20bio%20Lac-St-Jean/Pr%C3%A9sentation%20punaise%20terne%208%20mars%202012.pdf>

- **Thrips (*Frankliniella occidentalis/tritici*)**

Lemaire, É., 2011. Les thrips et le bronzage sur fraises: état des connaissances. MAPAQ, Direction régionale de la Capitale-Nationale.

Pearsall, I. A. and Myers, J. H., 2001. Spatial and temporal patterns of dispersal of western flower thrips (Thysanoptera : Thripidae) in nectarine orchards in British Columbia. *Journal of Economic Entomology*. 94, 4: 831-843

Pommier, J.-J., 1998. Thrips control using mist in strawberries. *Arboriculture fruitière*. 513, 25-28

Sampson, C., Kirk, W. 2016. Predatory mites double the economic injury level of *Frankliniella occidentalis* in strawberry, *BioControl* 61:661-669.

Shipp, J.L., Ward, K., Gillespie, T., 1996. Influence of temperature and vapor pressure deficit on the rate of predation by the predatory mite, *Amblyseius cucumeris*, on *Frankliniella occidentalis*. *Entomologia Experimentalis et Applicata* 78: 31-38.

Skirvin, D.J., Kravar-Garde L., Reynolds K., Jones, J., Mead, A., Fenlon, J., 2007. Supplemental food affects thrips predation and movement of *Orius laevigatus* (Hemiptera: Anthocoridae) and *Neoseiulus cucumeris* (Acari: Phytoseiidae). *Bulletin of Entomological Research*, volume 97, issue 3, pp. 309-315.

Steiner, M. Y. and Medhurst, A., 2003. Western flower thrips management strategies for strawberries. Report #BS00002. Horticulture Australia Limited, Sydney, Australia

- **Tétranyques à deux points (*Tetranychus urticae*)**

Caron, J., Laverdière, L., Roy, M., 2001. Mise au point finale et validation d'une stratégie de lutte intégrée contre le tétranyque à deux points dans les fraisières à production continue à l'aide du prédateur *Amblyseius fallacis*. Projet #204-08-981127

Fraulo, A., Liburd, O., 2007. Biological control of twospotted spider mite, *Tetranychus urticae*, with predatory mite, *Neoseiulus californicus*, in strawberries, Experimental and Applied Acarology, 43 :109-119

Rhodes, E., Liburd, O., 2005. A predatory mite, *Neoseiulus californicus* (McGregor). UF/IFAS Featured Creatures. EENY-359.

Rhodes, E., Liburd, O., Kelts, C., Rondon, S., Francis, R., 2006. Comparison of single and combination treatments of *Phytoseiulus persimilis*, *Neoseiulus californicus*, and Acramite (bifenazate) for control of twospotted spider mites in strawberries. Exp. Appl. Acarol. 39: 213-225.

Schausberger, P., Croft, B., 1999. Activity, feeding, and development among larvae of specialist and generalist phytoseiid mite species (Acari: Phytoseiidae). Environ. Entomol. 28: 322-329.

Wu, S., Xie, H., Li, M., Xu, X., Lei, Z. 2016. Highly virulent Beauvaria bassiana strains against the two-spotted spider mite, *Tetranychus urticae*, show no pathogenicity against five phytoseiid mite species, Exp Appl Acarol 70:421-435

- **Tarsonème (*Phytonemus pallidus*)**

Gobin, B., Audenaert, J., Vissers, M., Van Delsen, B., Vlaeminck, M. and Pauwels, E., 2013. Broad mite control in woody ornamentals: developing an integrated pest management solution. Acta Hortic. 990, 47-53

Svensson, B., 2009. Successful Bio-control of the strawberry mite *Phytonemus pallidus* with the predatory mite *Neoseiulus cucumeris* in organic outdoor production of strawberries (*Fragaria x ananassa* Duch.) in Sweden. Strawberry Symposium, Acta Hort. 842, HIS

Tremblay, J., Bouchard, A., Lafontaine, P., 2015. Évaluation du potentiel de certains insectes et acariens prédateurs pour le contrôle du tarsonème du fraisier, *Phytonemus pallidus* (Banks). Rapport final réalisé dans le cadre du programme Prime-Vert, sous-volet 11.1 . Appui à la Stratégie phytosanitaire québécoise en agriculture. Numéro du projet : CIEL-1-11-1581

Verschoor, J., Otma, E., Qiu, Y., Van Kruistum, G., Hoek, J., 2015. Controlled Atmosphere Temperature Treatment: Non-Chemical (Quarantine) Pest Control in Fresh Plant Products. Acta Hort. 1071, ISHS 2015; p. 253-58.

- **Anthracnose (*Colletotrichum* spp.)**

Dangovish, O., Bolda, M., Kaur, S., Mochizuki, M., Marcum, D., Epstein, L., 2012. Drip irrigation in California strawberry nurseries to reduce the incidence of *Colletotrichum acutatum* in fruit production. HortScience 47(3):368-373.

Debode J., Van Hemelrijck W., Baeyen S., Creemers P., Heungens K., Maes M., 2009. Quantitative detection and monitoring of *Colletotrichum acutatum* in strawberry leaves using real-time PCR. Plant Pathology 58, 504 -14.

Garrido C., Carbu M., Fernandez-Acero F. et al., 2009. Development of protocols for detection of *Colletotrichum acutatum* and monitoring of strawberry anthracnose using real-time PCR. Plant Pathology 58, 43 -51.

Martinez-Culebras P., Barrio E., Suarez-Fernandez M., Garcia-Lopez M., Querol A., 2002. RAPD analysis of *Colletotrichum* species isolated from strawberry and the design of specific primers for the identification of *C. fragariae*. Journal of Phytopathology 150, 680-6.

Martinez-Culebras P., Querol A., Suarez-Fernandez M., Garcia-Lopez M., Barrio E., 2003. Phylogenetic relationships among *Colletotrichum* pathogens of strawberry and design of PCR primers for their identification. Journal of Phytopathology 151, 135 -43.

Mertely, J., Martin, R., Peres, N.A., 2014. Control or root necrosis of strawberry caused by *Colletotrichum acutatum*, APS poster, University of Florida, Gulf Coast Research and Education Center. [https://strawberries.ces.ncsu.edu/wp-content/uploads/2014/09/APS-2014-poster-J-Mertely-root-necrosis.pdf?fwd=no](https://strawberries.ces.ncsu.edu/wp-content/uploads/2014/09/APS-2014-poster-J-Mertely-root-necrosis.pdf)

Merteley, J., Forcelini, B., Peres, N.A., nd. Root Necrosis of Strawberry Caused by *Colletotrichum acutatum*, University of Florida, IFAS Extension, <https://edis.ifas.ufl.edu/pdffiles/PP/PP12800.pdf>

Nam, M., Jeond, S., Lee, Y., Choi, J., Kim, G., 2006. Effects of nitrogen, phosphorus, potassium and calcium nutrition on strawberry anthracnose. Plant Pathology 55, 246-249.

Perez-Hernandez O., Nam M., Gleason M., Kim H., 2008. Development of a nested polymerase chain reaction assay for detection of *Colletotrichum acutatum* on symptomless strawberry leaves. Plant Disease 92, 1655 -61. Smith B, 2008. Epidemiol

Sreenivasaprasad S., Sharada K., Brown A., Mills P., 1996. PCR-based detection of *Colletotrichum acutatum* on strawberry. Plant Pathology 45, 650-5.

Walter, M., Braithwaite, B.; Smith, B. J.; Langford, G. I., 2008. Nutrient nitrogen management for disease control in strawberry. New Zealand Plant Protection 2008 Vol.61 pp.70-79 ref.8

- **Pourriture noire** (*Rhizoctonia fragariae*, *Pythium* spp., *Fusarium* sp., *Pratylenchus penetrans*), **pourriture du collet** (*Phytophthora cactorum*), **stèle rouge** (*Phytophthora fragariae*), **verticilliose** (*Verticillium dahliae*)

Berg, G., Fritze, A., Roskot, N., Smalla, K., 2001. Evaluation of potential biocontrol rhizobacteria from different host plants of *Verticillium dahliae*. Journal of Applied Microbiology 2001, 91, 963-971.

Coulombe, J., Langlois, D., Béclair, G., 2008. Les cultures de rotation et la biofumigation pour combattre les nématodes des lésions et la verticilliose dans le fraisier. Rapport final PSIH.

Demers, F., Hogue, R., Jeanne, T., 2014. Alternatives écologiques à la fumigation dans la culture de la fraise. Rapport final réalisé dans le cadre du programme Prime-Vert, sous-volet 11.1 . Appui à la Stratégie phytosanitaire québécoise en agriculture. numéro du projet : CPEM-2-11-1575

Elmer, W. H., LaMondia, J. A. 1999. Influence of ammonium sulfate and rotation crops on strawberry black root rot. Plant Dis. 83:119-123.

Landry, C., M. Marchand-Roy, J. Mainguy et M. Paradis., 2018. Développement d'une régie de culture misant sur l'amélioration de la santé des sols pour rétablir le potentiel de rendements de sites de fraisières en rangs nattés présentant un historique de déprérissement. Rapport final. IRDA. 30 pages.

Pinkerton, J. N., Ivors, K. L., Reeser, P. W., Bristow, P. R., Windom, G. E. 2002. The useof soil solarization for the management of soilborne plant pathogens in strawberry and red raspberry production. Plant Dis. 86:645-651.

Porras, M., Barrau, C., Arroyo, F. T., Santos, B., Blanco, C., and Romero, F.,2007. Reduction of *Phytophthora cactorum* in strawberry fields by *Trichoderma* spp. and soil solarization. Plant Dis. 91:142-146.

Subbarao, K. V., Kabir, Z., Martin, F. N., Koike, S. T., 2007. Management of soilborne diseases in strawberry using vegetable rotations. Plant Dis. 91:964-972.

Subbarao, K. V., Hubbard, J. C., and Koike, S.T.,1999. Evaluation of broccoli residue incorporation into field soil for *Verticillium* wilt control in cauliflower. Plant Dis. 83:124-129.